


## Universal Medication Schedule (UMS)



**Patients across all literacy levels can misunderstand medication directions**, although individuals with lower health literacy are more likely to misinterpret directions.

UMS uses specified timings such as ‘morning’, ‘noon’, ‘evening’, and ‘bedtime’ to create **explicit medication directions**.



Epic Systems has created a feature to translate standard directions to UMS directions.

### Standard Directions

Take one tablet by mouth twice daily  
Take one tablet by mouth every 12 hours  
Take one tablet by mouth at 9am and 9pm



### UMS Directions

Take 1 tablet by mouth in the morning and take 1 tablet in the evening

### UMS Directions...

Are identified as a best practice by the United States Pharmacopeia and the National Council for Prescription Drug Programs.

Increase understanding of when to take medication more than any other format, reducing errors

Simplify drug regimens

Improve communications about medications

Are a low resource strategy to improve adherence

### Medication Adherence...

Improved in patients who use UMS directions

Is key to managing chronic conditions

### Non-Adherence...

Affects **25-50%** of patients on prescription medication

Causes **125,000** annual deaths

Is the reason for **10%** of hospitalizations each year

Produces **\$100-\$289** billion annually in avoidable costs

Are you interested in adopting UMS at your organization or learning more?

Please contact Bhumi at Wisconsin Health Literacy, [bhumi@wisconsnliteracy.org](mailto:bhumi@wisconsnliteracy.org)

## References

- Bailey, S. C., Sarkar, U., Chen, A. H., Schillinger, D., & Wolf, M. S. (2012). Evaluation of language concordant, patient-centered drug label instructions. *Journal of General Internal Medicine*, 27(12), 1707–1713.  
<https://doi.org/10.1007/s11606-012-2035-3>
- Davis, T. C., Wolf, M. S., Bass, P. F., Thompson, J. A., Tilson, H. H., Neuberger, M., & Parker, R. M. (2006). Literacy and misunderstanding prescription drug labels. *Annals of Internal Medicine*, 145(12), 887.  
<https://doi.org/10.7326/0003-4819-145-12-200612190-00144>
- Iuga, A. & McGuire, M. (2014). Adherence and health care costs. *Risk Management and Healthcare Policy*, 35.  
<https://doi.org/10.2147/rmhp.s19801>
- Kebodeaux, C., Woodyard, J. L., Kachlic, M. D., Allen, S. M., Schadler, A. D., & Vouri, S. M. (2020). Student pharmacists' ability to organize complex medication regimens according to the universal medication schedule. *American Journal of Pharmaceutical Education*, 84(8). <https://doi.org/10.5688/ajpe7531>
- Viswanathan, M., Golin, C. E., Jones, C. D., Ashok, M., Blalock, S. J., Wines, R. C. M., Coker-Schwimmer, E. J. L., Rosen, D. L., Sista, P., & Lohr, K. N. (2012). Interventions to improve adherence to self-administered medications for chronic diseases in the United States. *Annals of Internal Medicine*, 157(11), 785.  
<https://doi.org/10.7326/0003-4819-157-11-201212040-00538>
- Wolf, M. S., Davis, T. C., Curtis, L. M., Bailey, S. C., Knox, J. A. P., Bergeron, A., Abbet, M., Shrank, W. H., Parker, R. M., & Wood, A. J. (2016). A patient-centered prescription drug label to promote appropriate medication use and adherence. *Journal of General Internal Medicine*, 31(12), 1482–1489. <https://doi.org/10.1007/s11606-016-3816-x>
- Wolf, M. S., Davis, T. C., Curtis, L. M., Webb, J. A., Bailey, S. C., Shrank, W. H., Lindquist, L., Ruo, B., Bocchini, M. V., Parker, R. M., & Wood, A. J. (2011). Effect of standardized, patient-centered label instructions to improve comprehension of prescription drug use. *Medical care*, 49(1), 96–100.  
<https://doi.org/10.1097/MLR.0b013e3181f38174>
- Wolf, M. S., Taitel, M. S., Jiang, J. Z., Curtis, L. M., Wismer, G. A., Wallia, A., & Parker, R. M. (2020). Prevalence of universal medication schedule prescribing and links to adherence. *American Journal of Health-System Pharmacy*, 77(3), 196–205. <https://doi.org/10.1093/ajhp/zxz305>